

Nonlinearity and Propagation Characteristics of Balanced Boolean Functions *

Jennifer Seberry
Xian-Mo Zhang
Yuliang Zheng

Department of Computer Science
The University of Wollongong
Wollongong, NSW 2522, AUSTRALIA
E-mail: jennie,xianmo,yuliang@cs.uow.edu.au

November 1993

*To appear in *Information and Computation*.

Running Head

Highly Nonlinear Balanced Boolean Functions

Contact Person

Dr Yuliang Zheng
Department of Computer Science
The University of Wollongong
Wollongong
NSW 2522
AUSTRALIA
E-mail: {jennie,xianmo,yuliang}@cs.uow.edu.au

Abstract

Three important criteria for cryptographically strong Boolean functions are balance, nonlinearity and the propagation criterion. The main contribution of this paper is to reveal a number of interesting properties of balance and nonlinearity, and to study systematic methods for constructing Boolean functions that satisfy some or all of the three criteria. We show that concatenating, splitting, modifying and multiplying (in the sense of Kronecker) sequences can yield balanced Boolean functions with a very high nonlinearity. In particular, we show that balanced Boolean functions obtained by modifying and multiplying sequences achieve a nonlinearity higher than that attainable by any previously known construction method. We also present methods for constructing balanced Boolean functions that are highly nonlinear and satisfy the strict avalanche criterion (SAC). Furthermore we present methods for constructing highly nonlinear balanced Boolean functions satisfying the propagation criterion with respect to *all but one or three* vectors. A technique is developed to transform the vectors where the propagation criterion is not satisfied in such a way that the functions constructed satisfy the propagation criterion of high degree while preserving the balance and nonlinearity of the functions. The algebraic degrees of functions constructed are also discussed.

Key Words

Bent Functions, Boolean Functions, Cryptography, Data Security, Hadamard Matrices, Nonlinearity, S-boxes, Sequences, Strict Avalanche Criterion.

1 Introduction

A Boolean function of n input coordinates is said to satisfy *the propagation criterion with respect to a non-zero vector* if complementing input coordinates according to the vector results in the output of the function being complemented 50% of the time over all possible input vectors, and to satisfy *the propagation criterion of degree k* if complementing k or less input coordinates results in the output of the function being complemented 50% of the time over all possible input vectors. Another important criterion is the strict avalanche criterion (SAC) that coincides with the propagation criterion of degree 1. The SAC was first introduced by Webster and Tavares (1985; 1986), was generalized in one direction by Forré (1989) and in another direction by Adams and Tavares (1990a). A combination of the two generalizations was studied in (Preneel *et al.*, 1991b; Preneel *et al.*, 1991a).

The nonlinearity of a Boolean function is defined as the minimum distance from the function to the affine functions. A cryptosystem that employs functions with a low nonlinearity is vulnerable to many cryptanalytic attacks, including the linear cryptanalysis discovered by Matsui (1994). It is well known that bent functions possess the highest nonlinearity and satisfy the propagation criterion with respect to *all* non-zero vectors (Dillon, 1972). However two drawbacks of bent functions prohibit their direct application in practice. The first drawback is that they are not balanced, and the second drawback is that they exist only when the number of input coordinates is even. Cryptographic applications, such as the design of strong substitution boxes (S-boxes), often require that when input coordinates of a Boolean function are selected independently, at random, the output of the function must behave as a uniformly distributed random variable (Kam & Davida, 1979; Adams & Tavares, 1990a; Seberry *et al.*, 1993). In other words, the function has to be balanced. Some practical applications need Boolean functions with an odd number of input coordinates.

This paper studies properties and constructions of nonlinear, balanced functions. We present a number of methods for constructing highly nonlinear balanced functions. These include concatenating, splitting, modifying and multiplying (in the sense of Kronecker) sequences. It is interesting

to note that balanced functions obtained by modifying and multiplying sequences achieve a nonlinearity higher than that attainable by any previously known construction method. We also present methods for systematically constructing balanced functions satisfying the SAC. When $n = 2k + 1$, where n is the number of input coordinates, the nonlinearity of functions constructed is at least $2^{2k} - 2^k$, and when $n = 2k$, it is at least $2^{2k-1} - 2^k$.

Furthermore we present methods for constructing balanced functions satisfying high degree propagation criterion. More precisely, when $n = 2k + 1$, we construct nonlinearly balanced functions that satisfy the propagation criterion with respect to *all but one* non-zero vectors, and when $n = 2k$, functions we construct are balanced and also satisfy the propagation criterion with respect to *all but three* non-zero vectors. We also show that the vectors where the propagation criterion is not satisfied can be transformed into other vectors. As a consequence, we obtain balanced functions satisfying the propagation criterion of degree $2k$ when $n = 2k + 1$, and balanced functions satisfying the propagation criterion of degree $\frac{4k}{3}$ when $n = 2k$. The nonlinearity of functions constructed is at least $2^{2k} - 2^k$ when $n = 2k + 1$, and $2^{2k-1} - 2^k$ when $n = 2k$.

The organization of the remainder of the paper is as follows: in Section 2 we introduce notations and definitions used in this paper. In Section 3 we prove results on the nonlinearity and balance of functions including those obtained by concatenating or splitting bent sequences. In Section 4, we show methods for constructing highly nonlinear balanced functions by modifying and multiplying sequences. Our construction methods for highly nonlinear balanced functions satisfying the SAC are presented in Section 5, while methods for highly nonlinear balanced functions satisfying high degree propagation criterion are presented in Section 6. The paper is closed by a discussion of future work in Section 7.

2 Preliminaries

We consider functions from V_n to $GF(2)$ (or simply functions on V_n), where V_n is the vector space of n tuples of elements from $GF(2)$. These functions are also called Boolean functions. Note that functions on V_n can be represented by polynomials of n coordinates. We are particularly interested in the *algebraic normal form* representation in which a function is viewed as the sum of products of coordinates. The *algebraic degree* of a function is the number of coordinates in the longest product when the function is represented in the algebraic normal form. To distinguish between a vector of coordinates and an individual coordinate, the former will be strictly denoted by x, y or z , while the latter strictly by x_i, y_i, z_i, u or v , where i is an index.

Let f be a function on V_n . The $(1, -1)$ -sequence defined by $((-1)^{f(\alpha_0)}, (-1)^{f(\alpha_1)}, \dots, (-1)^{f(\alpha_{2^n-1})})$ is called the *sequence* of f , and the $(0, 1)$ -sequence defined by $(f(\alpha_0), f(\alpha_1), \dots, f(\alpha_{2^n-1}))$ is called the *truth table* of f , where $\alpha_i, 0 \leq i \leq 2^n - 1$, denotes the vector in V_n whose integer representation is i . A $(0, 1)$ -sequence ($(1, -1)$ -sequence) is said to be *balanced* if it contains an equal number of zeros and ones (ones and minus ones). A function is balanced if its sequence is balanced.

Obviously if (a_0, \dots, a_{2^n-1}) and (b_0, \dots, b_{2^n-1}) are the sequences of functions f_1 and f_2 on V_n respectively, then $(a_0 b_0, \dots, a_{2^n-1} b_{2^n-1})$ is the sequence of $f(x) \oplus g(x)$, where $x = (x_1, x_2, \dots, x_n)$. In particular, $-(a_0, \dots, a_{2^n-1}) = (-a_0, \dots, -a_{2^n-1})$ is the sequence of $1 \oplus f_1(x)$.

An *affine* function f on V_n is a function that takes the form of $f(x) = a_1 x_1 \oplus \dots \oplus a_n x_n \oplus c$, where $a_j, c \in GF(2), j = 1, 2, \dots, n$. Furthermore f is called a *linear* function if $c = 0$. The sequence of an affine (or linear) function is called an *affine (or linear) sequence*. The *Hamming weight* of a $(0, 1)$ -sequence (or vector) α , denoted by $W(\alpha)$, is the number of ones in α . The *Hamming distance* between two sequences α and β of the same length, denoted by $d(\alpha, \beta)$, is the number of positions where the two sequences differ. Given two functions f and g on V_n , the Hamming distance between

them is defined as $d(f, g) = d(\xi_f, \xi_g)$, where ξ_f and ξ_g are the truth tables of f and g respectively. The *nonlinearity* of f , denoted by N_f , is the minimal Hamming distance between f and all affine functions on V_n , i.e., $N_f = \min_{i=0,1,\dots,2^{n+1}-1} d(f, \varphi_i)$ where $\varphi_0, \varphi_1, \dots, \varphi_{2^{n+1}-1}$ denote the affine functions on V_n .

The following notation will be used in this paper. Let $\alpha = (a_1, \dots, a_n)$ and $\beta = (b_1, \dots, b_n)$ be two sequences (or vectors), the *scalar product* of α and β , denoted by $\langle \alpha, \beta \rangle$, is defined as the sum of the component-wise multiplications. In particular, when α and β are from V_n , $\langle \alpha, \beta \rangle = a_1 b_1 \oplus \dots \oplus a_n b_n$, where the addition and the multiplication are over $GF(2)$, and when α and β are $(1, -1)$ -sequences, $\langle \alpha, \beta \rangle = a_1 b_1 + \dots + a_n b_n$, where the addition and the multiplication are over the reals.

The *Kronecker product* of an $m \times n$ matrix A and an $s \times t$ matrix B , denoted by $A \otimes B$, is an $ms \times nt$ matrix defined by

$$A \otimes B = \begin{bmatrix} a_{11}B & a_{12}B & \cdots & a_{1n}B \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1}B & a_{m2}B & \cdots & a_{mn}B \end{bmatrix}$$

where a_{ij} is the element in the i th row and the j th column of A . In particular, the Kronecker product of a sequence α of length m and a sequence β of length n is a sequence of length mn defined by $\alpha \otimes \beta = (a_1\beta, a_2\beta, \dots, a_m\beta)$, where a_i is the i th element in α .

A $(1, -1)$ -matrix H of order n is called a *Hadamard matrix* if $HH^t = nI_n$, where H^t is the transpose of H and I_n is the identity matrix of order n . It is well known that the order of a Hadamard matrix is 1, 2 or divisible by 4 (Wallis *et al.*, 1972). A special kind of Hadamard matrix, called *Sylvester-Hadamard matrix* or *Walsh-Hadamard matrix*, will be relevant to this paper. A Sylvester-Hadamard matrix of order 2^n , denoted by H_n , is generated by the following recursive relation

$$H_0 = 1, H_n = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \otimes H_{n-1}, n = 1, 2, \dots$$

Note that H_n can be represented as $H_n = H_s \otimes H_t$ for any s and t with $s + t = n$. such matrices are closely related to linear functions, as is shown in the following well-known lemma.

Lemma 1 Write $H_n = \begin{bmatrix} \ell_0 \\ \ell_1 \\ \vdots \\ \ell_{2^n-1} \end{bmatrix}$ where ℓ_i is a row of H_n . Then ℓ_i is the sequence of $h_i = \langle \alpha_i, x \rangle$,

a linear function, where α_i is a vector in V_n whose integer representation is i and $x = (x_1, \dots, x_n)$. Conversely the sequence of any linear function on V_n is a row of H_n .

From Lemma 1 the rows of H_n comprise the sequences of all linear functions on V_n . Consequently the rows of $\pm H_n$ comprise the sequences of all *affine* functions on V_n .

The following notation is very useful in obtaining the functional representation of a concatenated sequence. Let $\delta = (i_1, i_2, \dots, i_p)$ be a vector in V_p . Then D_δ is a function on V_p defined by

$$D_\delta(y_1, y_2, \dots, y_p) = (y_1 \oplus i_1 \oplus 1) \cdots (y_p \oplus i_p \oplus 1).$$

Using this notation one can readily prove

Lemma 2 Let $f_0, f_1, \dots, f_{2^p-1}$ be functions on V_q . Let ξ_i the sequence of $f_i, i = 0, 1, \dots, 2^p - 1$, and let ξ be the concatenation of $\xi_0, \xi_1, \dots, \xi_{2^p-1}$, namely, $\xi = (\xi_0, \xi_1, \dots, \xi_{2^p-1})$. Then ξ is the sequence of the following function on V_{p+q}

$$f(y, x) = \bigoplus_{i=0}^{2^p-1} D_{\alpha_i}(y) f_i(x)$$

where $y = (y_1, \dots, y_p), x = (x_1, \dots, x_q)$ and α_i is the vector in V_p whose integer representation is i .

For instance, if ξ_1, ξ_2 are the sequences of functions f_1, f_2 on V_n , then $\eta = (\xi_1, \xi_2)$ is the sequence of $(1 \oplus u)f_1(x_1, \dots, x_n) \oplus uf_2(x_1, \dots, x_n)$, a function on V_{n+1} .

We now introduce the concept of bent functions.

Definition 1 A function f on V_n is called a bent function if

$$2^{-\frac{n}{2}} \sum_{x \in V_n} (-1)^{f(x) \oplus \langle \beta, x \rangle} = \pm 1$$

for all $\beta \in V_n$. Here $f(x) \oplus \langle \beta, x \rangle$ is regarded as a real-valued function. The sequence of a bent function is called a bent sequence.

From the definition we can see that bent functions on V_n exist only when n is even. It was Rothaus who first introduced and studied bent functions in the 1960s, although his pioneering work was not published in the open literature until some ten years later (Rothaus, 1976). Other issues related to bent functions, such as properties, constructions and counting, can be found in (Adams & Tavares, 1990a; Kumar & Scholtz, 1983; Lempel & Cohn, 1982; Olsen *et al.*, 1982; Yarlagadda & Hershey, 1989). Kumar, Scholtz and Welch (1985) defined and studied bent functions from Z_q^n to Z_q , where q is a positive integer. Applications of bent functions to digital communications, coding theory and cryptography can be found in such as (Adams & Tavares, 1990b; Detombe & Tavares, 1993; Lempel & Cohn, 1982; Losev, 1987; MacWilliams & Sloane, 1978; Meier & Staffelbach, 1990; Nyberg, 1991; Olsen *et al.*, 1982; Seberry *et al.*, 1993).

The following result can be found in an excellent survey of bent functions by Dillon (1972).

Lemma 3 Let f be a function on V_n , and let ξ be the sequence of f . Then the following four statements are equivalent:

(i) f is bent.

(ii) $\langle \xi, \ell \rangle = \pm 2^{\frac{1}{2}n}$ for any affine sequence ℓ of length 2^n .

(iii) $f(x) \oplus f(x \oplus \alpha)$ is balanced for any non-zero vector $\alpha \in V_n$.

(iv) $f(x) \oplus \langle \alpha, x \rangle$ assumes the value one $2^{n-1} \pm 2^{\frac{1}{2}n-1}$ times for any $\alpha \in V_n$.

By (iv) of Lemma 3, if f is a bent function on V_n , then $f(x) \oplus h(x)$ is also a bent function for any affine function h on V_n . This property will be employed in constructing highly nonlinear balanced functions to be described in Sections 5 and 6.

The notion of *strict avalanche criterion (SAC)* was first introduced by Webster and Tavares (1985; 1986).

Definition 2 A function f on V_n is said to satisfy the SAC if complementing any single input coordinate results in the output of f being complemented half the times over all input vectors, namely, $f(x) \oplus f(x \oplus \alpha)$ is a balanced function for any vector $\alpha \in V_n$ whose Hamming weight is 1.

In this paper we are also concerned with the propagation criterion, which was introduced in (Adams & Tavares, 1990a; Preneel *et al.*, 1991b) as a generalization of the SAC.

Definition 3 Let f be a function on V_n . We say that f satisfies

1. the propagation criterion with respect to a non-zero vector α in V_n if $f(x) \oplus f(x \oplus \alpha)$ is a balanced function.
2. the propagation criterion of degree k if it satisfies the propagation criterion with respect to all $\alpha \in V_n$ with $1 \leq W(\alpha) \leq k$.

Note that the SAC is equivalent to the propagation criterion of degree 1, and that the *perfect nonlinearity* studied by Meier and Staffelbach (1990) is equivalent to the propagation criterion of degree n .

Now it becomes clear that when n is even, only bent functions fulfill the propagation criterion of the maximal degree n . Another property of bent functions is that they possess the highest possible nonlinearity. This will be discussed in more detail in the next section.

3 Properties of Balance and Nonlinearity

This section presents a number of results related to balance and nonlinearity. These include upper bounds for nonlinearity and properties of concatenated and split sequences.

3.1 Upper Bounds on Nonlinearity

It is well-known that the maximum nonlinearity of functions on V_n coincides with the covering radius of the first order binary Reed-Muller code $R(1, n)$ of length 2^n (MacWilliams & Sloane, 1978). By translating an upper bound on the covering radius of $R(1, n)$ (Cohen *et al.*, 1985), we have:

Lemma 4 For any function f on V_n , the nonlinearity N_f of f satisfies $N_f \leq 2^{n-1} - 2^{\frac{1}{2}n-1}$.

Remark 1 A function on V_n attains the upper bound for nonlinearities, $2^{n-1} - 2^{\frac{1}{2}n-1}$, if and only if it is bent.

Recall that bent functions are not balanced. From Remark 1, balanced functions can not attain the upper bound for nonlinearities, namely $2^{n-1} - 2^{\frac{1}{2}n-1}$. A slightly improved upper bound for the nonlinearities of balanced functions can be obtained by noting the fact that a balanced function assumes the value one an even number of times.

Lemma 5 Let f be a balanced function on V_n ($n \geq 3$). Then the nonlinearity N_f of f is given by

$$N_f \leq \begin{cases} 2^{n-1} - 2^{\frac{1}{2}n-1} - 2, & n \text{ even} \\ \lfloor \lfloor 2^{n-1} - 2^{\frac{1}{2}n-1} \rfloor \rfloor, & n \text{ odd} \end{cases}$$

where $\lfloor \lfloor x \rfloor \rfloor$ denotes the maximum even integer less than or equal to x .

Proof. Note that the length of the sequence of a function is even. Also note that the truth table of f contains an even number of ones and that all affine sequences contain an even number of ones. Then $N_f = \min_{i=0,1,\dots,2^{n+1}-1} d(f, \varphi_i)$, where $\varphi_0, \varphi_1, \dots, \varphi_{2^{n+1}-1}$ denote the affine functions on V_n , must be even. On the other hand, since f is not bent, by Remark 1 we have $N_f < 2^{n-1} - 2^{\frac{1}{2}n-1}$. This proves the lemma. \square

For V_2 , there are six balanced sequences, namely

$$\pm(1, 1, -1, -1), \pm(1, -1, 1, -1), \pm(1, -1, -1, 1)$$

all of which are linear. Therefore there are no nonlinearly balanced functions on V_2 .

3.2 Concatenating Sequences

First we establish a lemma that is very useful in calculating the nonlinearity of a function.

Lemma 6 *Let f and g be functions on V_n whose sequences are ξ_f and ξ_g respectively. Then the distance between f and g can be calculated by $d(f, g) = 2^{n-1} - \frac{1}{2}\langle \xi_f, \xi_g \rangle$.*

Proof. $\langle \xi_f, \xi_g \rangle = \sum_{f(x)=g(x)} 1 - \sum_{f(x) \neq g(x)} 1 = 2^n - 2 \sum_{f(x) \neq g(x)} 1 = 2^n - 2d(f, g)$. This proves the lemma. \square

The following lemma gives a lower bound on the nonlinearity of a function obtained by concatenating the sequences of two functions.

Lemma 7 *Let f_1 and f_2 be functions on V_n , and let g be a function on V_{n+1} defined by*

$$g(u, x_1, \dots, x_n) = (1 \oplus u)f_1(x_1, \dots, x_n) \oplus uf_2(x_1, \dots, x_n). \quad (1)$$

Suppose that ξ_1 and ξ_2 , the sequences of f_1 and f_2 respectively, satisfy $\langle \xi_1, \ell \rangle \leq P_1$ and $\langle \xi_2, \ell \rangle \leq P_2$ for any affine sequence ℓ of length 2^n , where P_1 and P_2 are positive integers. Then the nonlinearity of g satisfies $N_g \geq 2^n - \frac{1}{2}(P_1 + P_2)$.

Proof. Note that $\xi = (\xi_1, \xi_2)$ is the sequence of g . Let ψ be an arbitrary affine function on V_{n+1} and let L be the sequence of ψ . Then L must take the form of $L = (\ell, \pm\ell)$ where ℓ is an affine sequence of length 2^n . Note that $\langle \xi, L \rangle = \langle \xi_1, \ell \rangle \pm \langle \xi_2, \ell \rangle$ and thus $|\langle \xi, L \rangle| \leq P_1 + P_2$. By Lemma 6, we have $d(g, \psi) = 2^n - \frac{1}{2}\langle \xi, L \rangle \geq 2^n - \frac{1}{2}(P_1 + P_2)$. Since ψ is arbitrary, we have $N_g \geq 2^n - \frac{1}{2}(P_1 + P_2)$, and this completes the proof. \square

As bent functions do not exist on V_{2k+1} , an interesting question is what functions on V_{2k+1} are highly nonlinear. The following result, as a special case of Lemma 7, shows that such functions can be obtained by concatenating bent sequences. This construction has been discovered by Meier and Staffelbach in (1990).

Corollary 1 *In the construction (1), if both f_1 and f_2 are bent functions on V_{2k} , then $N_g \geq 2^{2k} - 2^k$.*

A similar result can be obtained when the sequences of four functions are concatenated.

Lemma 8 Let f_0, f_1, f_2 and f_3 be functions on V_n whose sequences are ξ_0, ξ_1, ξ_2 and ξ_3 respectively. Assume that $\langle \xi_i, \ell \rangle \leq P_i$ for each $0 \leq i \leq 3$ and for each affine sequence ℓ of length 2^n , where each P_i is a positive integer. Let g be a function on V_{n+2} defined by

$$g(y, x) = \bigoplus_{i=0}^3 D_{\alpha_i}(y) f_i(x) \quad (2)$$

where $y = (y_1, y_2)$, $x = (x_1, \dots, x_n)$ and α_i is a vector in V_2 whose integer representation is i . Then $N_g \geq 2^{n+1} - \frac{1}{2}(P_0 + P_1 + P_2 + P_3)$. In particular, when n is even and f_0, f_1, f_2 and f_3 are all bent functions on V_n , $N_g \geq 2^{n+1} - 2^{\frac{1}{2}n+1}$.

The proof for Lemma 8 is similar to that for Lemma 7, and hence is omitted. It is a simple exercise to further generalize the lemma to the case where the sequences of 2^t , $t \geq 1$, functions are concatenated.

By selecting proper starting functions in (1) and (2), the resulting functions can be balanced. For instance, in (1), if both f_1 and f_2 are balanced, or the number of times f_1 assumes the value one is equal to that f_2 assumes the value zero, the resulting function g is balanced.

3.3 Splitting Sequences

We have discussed the concatenation of sequences of functions including bent functions. The following lemma deals with the other direction, namely splitting bent sequences.

Lemma 9 Let $f(x_1, x_2, \dots, x_{2k})$ be a bent function on V_{2k} , η_0 be the sequence of $f(0, x_2, \dots, x_{2k})$, and η_1 be the sequence of $f(1, x_2, \dots, x_{2k})$. Then for any affine sequence ℓ of length 2^{2k-1} , we have $-2^k \leq \langle \eta_0, \ell \rangle \leq 2^k$ and $-2^k \leq \langle \eta_1, \ell \rangle \leq 2^k$.

Proof. We only give a proof for $-2^k \leq \langle \eta_0, \ell \rangle \leq 2^k$. The other half can be proved in the same way. Since $f(x_1, x_2, \dots, x_{2k}) = (1 \oplus x_1)f(0, x_2, \dots, x_{2k}) \oplus x_1 f(1, x_2, \dots, x_{2k})$, $\eta = (\eta_0, \eta_1)$ is the sequence of $f(x_1, x_2, \dots, x_{2k})$. Let $L = (\ell, \ell)$ and $L' = (\ell, -\ell)$. By Lemma 1, both L and L' are affine sequences of length 2^{2k} .

Suppose that $-2^k \leq \langle \eta_0, \ell \rangle \leq 2^k$ is not true. Without loss of generality assume that $\langle \eta_0, \ell \rangle > 2^k$. There are two cases that have to be considered: $\langle \eta_1, \ell \rangle > 0$ and $\langle \eta_1, \ell \rangle < 0$. In the first case we have $\langle \eta, L \rangle \geq \langle \eta_0, \ell \rangle + \langle \eta_1, \ell \rangle > 2^k$, and in the second case we have $\langle \eta, L' \rangle \geq \langle \eta_0, \ell \rangle + \langle \eta_1, -\ell \rangle = \langle \eta_0, \ell \rangle + (-1)\langle \eta_1, \ell \rangle > 2^k$, both of which contradict the fact that $\langle \eta, L \rangle = \pm 2^k$ (see also (ii) of Lemma 3). This completes the proof. \square

A consequence of Lemma 9 is that the nonlinearity of split functions $f(0, x_2, \dots, x_{2k})$ and $f(1, x_2, \dots, x_{2k})$ is at least $2^{2k-2} - 2^{k-1}$. It is interesting to note that concatenating and splitting bent sequences both achieve the same nonlinearity.

Splitting bent sequences can also result in balanced functions. Let ℓ_i be the i th row of H_k where $i = 0, 1, \dots, 2^k - 1$. Note that ℓ_0 is an all-one sequence while $\ell_1, \ell_2, \dots, \ell_{2^k-1}$ are all balanced sequences. The concatenation of the rows, $(\ell_0, \ell_1, \dots, \ell_{2^k-1})$, is a bent sequence (Adams & Tavares, 1990a). Denote by $f(x_1, x_2, \dots, x_{2k})$ the function corresponding to the bent sequence. Let ξ be the second half of the bent sequence, namely, $\xi = (\ell_{2^k-1}, \ell_{2^k-1+1}, \dots, \ell_{2^k-1})$. Then ξ is the sequence of $f(1, x_2, \dots, x_{2k})$. Since all ℓ_i , $i = 2^{k-1}, 2^{k-1} + 1, \dots, 2^k - 1$, are balanced, $f(1, x_2, \dots, x_{2k})$ is a balanced function. The nonlinearity of the function is at least $2^{2k-2} - 2^{k-1}$.

By permuting $\{\ell_{2^k-1}, \ell_{2^k-1+1}, \dots, \ell_{2^k-1}\}$, we obtain a new balanced sequence

$$\xi' = (\ell'_{2^k-1}, \ell'_{2^k-1+1}, \dots, \ell'_{2^k-1})$$

that has the same nonlinearity. Now let

$$\xi'' = (e_{2^{k-1}} \ell'_{2^{k-1}}, e_{2^{k-1}+1} \ell'_{2^{k-1}+1}, \dots, e_{2^k-1} \ell'_{2^k-1}),$$

where each e_i is independently selected from $\{1, -1\}$. ξ'' is also a balanced sequence with the same nonlinearity. The total number of different balanced sequences obtained by permuting and changing signs is $2^{2^{k-1}} \cdot 2^{k-1}!$.

3.4 An Invariance Property

Next we examine properties of functions with respect to the affine transformation of coordinates. Let f be a function on V_n , A a nondegenerate matrix of order n with entries from $GF(2)$, and b a vector in V_n . Then $f^*(x) = f(xA \oplus b)$ defines a new function on V_n , where $x = (x_1, x_2, \dots, x_n)$. It is obvious that the algebraic degree of f^* is the same as that of f .

On the other hand, since A is nondegenerate, $xA \oplus b$ is an one-to-one mapping on V_n . Hence the truth table of f^* contains exactly the same number of ones as that of f . This indicates that the balance of a function is preserved under the affine transformation of coordinates.

Now let φ be an affine function on V_n and let $\varphi^*(x) = \varphi(xA \oplus b)$. It is easy to verify that $d(f, \varphi) = d(f^*, \varphi^*)$. Since A is nondegenerate, φ^* will run through all affine functions on V_n while φ runs through all affine functions on V_n . This proves that the nonlinearity of f^* is the same as that of f (Meier & Staffelbach, 1990).

Finally we consider the propagation characteristics under the affine transformation of coordinates. Let α be a nonzero vector in V_n . $f^*(x) \oplus f^*(x \oplus \alpha)$ is balanced if and only if

$$\begin{aligned} f(xA \oplus b) \oplus f((x \oplus \alpha)A \oplus b) &= f(xA \oplus b) \oplus f((xA \oplus b) \oplus \alpha A) \\ &= f(y) \oplus f(y \oplus \beta) \end{aligned}$$

is balanced, where $y = xA \oplus b$ and $\beta = \alpha A$. Since A is nondegenerate and α is a nonzero vector, β is a nonzero vector. In addition, $y = xA \oplus b$ will run through V_n while x runs through V_n . Therefore the number of vectors in V_n where the propagation criterion is satisfied remains unchanged under the affine transformation. To summarize the discussions, we have

Lemma 10 *The algebraic degree, the Hamming weight of the truth table, the nonlinearity, and the number of vectors with respect to which the propagation criterion is satisfied, of a function are invariant under the affine transformation of coordinates.*

4 Highly Nonlinear Balanced Functions

Recall that a bent sequence of length 2^{2k} contains $2^{2k-1} + 2^{k-1}$ ones and $2^{2k-1} - 2^{k-1}$ zeros, or vice versa. As is observed by Meier and Staffelbach (1990), complementing 2^{k-1} positions in a bent sequence yields a balanced function on V_{2k} having a nonlinearity of at least $2^{2k-1} - 2^k$. This nonlinearity is the same as that obtained by concatenating four bent sequences of length 2^{2k-2} (see Lemma 8). We note, however, that concatenation is superior to complementation in that it is far easier to discuss cryptographic properties such as the propagation characteristics of functions obtained by concatenation than to discuss properties by complementation.

Now we consider the case of V_{2k+1} . As the maximum nonlinearity of functions on V_n coincides with the covering radius of the first order binary Reed-Muller code $R(1, n)$ of length 2^n , using a result of (Patterson & Wiedemann, 1983), we can construct *unbalanced* functions on V_{2k+1} , $k \geq 7$, whose nonlinearity is at least $2^{2k} - \frac{108}{128}2^k$, a higher value than $2^{2k} - 2^k$ achieved by the construction

in Corollary 1. One might be tempted to think that modifying the sequences in (Patterson & Wiedemann, 1983) would result in balanced functions with a higher nonlinearity than that obtained by concatenating or splitting bent sequences. We find that it is not the case. We take V_{15} for an example. The Hamming weight of the sequences on V_{15} , which have the largest nonlinearity of 16276, is 16492. Changing 54 positions makes them balanced. The nonlinearity of the resulting functions is 16222, smaller than 16256 achieved by concatenating two bent sequences of length 2^{14} (see Corollary 1).

To summarize the above discussions, so far the best result on constructing nonlinearly balanced functions on V_{2k} is by concatenating four bent sequences of length 2^{2k-2} , while the best result on V_{2k+1} is by concatenating two bent sequences of length 2^{2k} , or by splitting a bent sequence of length 2^{2k+2} .

In the following we show how to modify bent sequences of length 2^{2k} constructed from Hadamard matrices in such a way that the resulting functions on V_{2k} are balanced and have a much higher nonlinearity than that attainable by concatenating four bent sequences. This result, in conjunction with sequences in (Patterson & Wiedemann, 1983), allows us to construct balanced functions on V_{2k+1} , $k \geq 14$, that have a higher nonlinearity than that achieved by concatenating or splitting bent sequences. These results represent a significant improvement to the previously known construction methods.

4.1 On V_{2k}

Note that an even number $n \geq 4$ can be expressed as $n = 4t$ or $n = 4t + 2$, where $t \geq 1$. As the first step towards our goal, we prove

Lemma 11 *For any integer $t \geq 1$ there exists*

- (i) *a balanced function f on V_{4t} such that $N_f \geq 2^{4t-1} - 2^{2t-1} - 2^t$,*
- (ii) *a balanced function f on V_{4t+2} such that $N_f \geq 2^{4t+1} - 2^{2t} - 2^t$.*

Proof. (i) Let ℓ_i be the i th row of H_{2t} where $i = 0, 1, \dots, 2^{2t} - 1$. Then $\xi = (\ell_0, \ell_1, \dots, \ell_{2^{2t}-1})$ is a bent sequence of length 2^{4t} .

Note that except for $\ell_0 = (1, 1, \dots, 1)$, all other ℓ_i ($i = 1, \dots, 2^{2t} - 1$) are balanced sequences of length 2^{2t} . Therefore replacing the all-one (or “flat”) leading sequence ℓ_0 with a balanced sequence renders ξ balanced. The crucial idea here is to select a replacement with a high nonlinearity, since the nonlinearity of the resulting function depends largely on that of the replacement.

The replacement we select is $\ell_0^* = (e_1, e_1, e_2, \dots, e_{2^{2t}-1})$, where e_i is the i th row of H_t . Note that the leading sequence in ℓ_0^* is e_1 but not $e_0 = (1, 1, \dots, 1)$. ℓ_0^* is a balanced sequence of length 2^{2t} , since all e_i , $i = 1, \dots, 2^{2t} - 1$, are balanced sequences of length 2^t . Replacing ℓ_0 by ℓ_0^* , we get a balanced sequence $\xi^* = (\ell_0^*, \ell_1, \dots, \ell_{2^{2t}-1})$.

Denote by f^* the function corresponding to the sequence ξ^* , and consider the nonlinearity of f^* . Let φ be an arbitrary affine function on V_{4t} , and let L be the sequence of φ . By Lemma 1, L is a row of $\pm H_{4t}$. Since $H_{4t} = H_{2t} \otimes H_{2t}$, L can be expressed as $L = \pm \ell_i \otimes \ell_j$, where ℓ_i and ℓ_j are two rows of H_{2t} . Assume that $\ell_i = (a_0, a_1, \dots, a_{2^{2t}-1})$. Then $L = \pm(a_0 \ell_j, a_1 \ell_j, \dots, a_{2^{2t}-1} \ell_j)$. A property of a Hadamard matrix is that its rows are mutually orthogonal. Hence $\langle \ell_p, \ell_q \rangle = 0$ for $p \neq q$. Thus

$$|\langle \xi^*, L \rangle| \leq |\langle \ell_0^*, \ell_j \rangle| + |\langle \ell_j, \ell_j \rangle| \leq |\langle \ell_0^*, \ell_j \rangle| + 2^{2t}.$$

We proceed to estimate $|\langle \ell_0^*, \ell_j \rangle|$. Note that $H_{2t} = H_t \otimes H_t$, ℓ_j can be expressed as $\ell_j = e_u \otimes e_v$, where e_u and e_v are rows of H_t . Write $e_u = (b_0, \dots, b_{2^t-1})$. Then $\ell_j = (b_0 e_v, \dots, b_{2^t-1} e_v)$. Similarly to the discussion for $|\langle \xi^*, L \rangle|$, we have

$$|\langle \ell_0^*, \ell_j \rangle| \leq \begin{cases} 2|\langle e_2, e_2 \rangle| = 2^{t+1}, & \text{if } v = 1, \\ |\langle e_v, e_v \rangle| = 2^t, & \text{if } v = 2, \dots, 2^t - 1, \\ 0, & \text{if } v = 0. \end{cases}$$

Thus $|\langle \ell_0^*, \ell_j \rangle| \leq 2^{t+1}$ and hence $|\langle \xi^*, L \rangle| \leq 2^{t+1} + 2^{2t}$.

By Lemma 6, $d(f^*, \varphi) \geq 2^{4t-1} - \frac{1}{2}|\langle \xi^*, L \rangle| \geq 2^{4t-1} - 2^{2t-1} - 2^t$. Since φ is arbitrary, $N_{f^*} \geq 2^{4t-1} - 2^{2t-1} - 2^t$.

(ii) Now consider the case of V_{4t+2} . Let ℓ_i , $i = 0, 1, \dots, 2^{2t+1} - 1$, be the i th row of H_{2t+1} . Then $\xi = (\ell_0, \ell_1, \dots, \ell_{2^{2t+1}-1})$ is a bent sequence of length 2^{4t+2} .

The replacement for the all-one leading sequence $\ell_0 = (1, 1, \dots, 1) \in V_{2t+1}$ is the following balanced sequence $\ell_0^* = (e_{2^t}, e_{2^t+1}, \dots, e_{2^{2t+1}-1})$, the concatenation of the 2^t th, the $(2^t + 1)$ th, \dots , and the $(2^{2t+1} - 1)$ th rows of H_{t+1} . Let $\xi^* = (\ell_0^*, \ell_1, \dots, \ell_{2^{2t+1}-1})$, and let f^* be the function corresponding to the balanced sequence.

Similarly to the case of V_{4t} , let φ be a affine function on V_{4t+2} and let L be its sequence. L can be expressed as $L = \pm \ell_i \otimes \ell_j$ where ℓ_i and ℓ_j are rows of H_{2t+1} . Hence

$$|\langle \xi^*, L \rangle| \leq |\langle \ell_0^*, \ell_j \rangle| + |\langle \ell_j, \ell_j \rangle| \leq |\langle \ell_0^*, \ell_j \rangle| + 2^{2t+1}.$$

Since ℓ_0^* is obtained by splitting the bent sequence $(e_0, e_1, \dots, e_{2^{2t+1}-1})$, where e_i is a row of H_{t+1} , by Lemma 9, we have $|\langle \ell_0^*, \ell_j \rangle| \leq 2^{t+1}$. From this it follows that $|\langle \xi^*, L \rangle| \leq 2^{t+1} + 2^{2t+1}$ and $N_{f^*} \geq 2^{4t+1} - 2^{2t} - 2^t$. \square

With the above result as a basis, we consider an iterative procedure to further improve the nonlinearity of a balanced function. Note that an even number $n \geq 4$ can be expressed as $n = 2^m$, $m \geq 2$, or $n = 2^s(2t + 1)$, $s \geq 1$ and $t \geq 1$.

Consider the case when $n = 2^m$, $m \geq 2$. We start with the bent sequence obtained by concatenating the rows of $H_{2^{m-1}}$. The sequence consists of $2^{2^{m-1}}$ sequences of length $2^{2^{m-1}}$. Now we replace the all-one leading sequence with a bent sequence of the same length, which is obtained by concatenating the rows of $H_{2^{m-2}}$. The length of the new leading sequence becomes $2^{2^{m-2}}$. It is replaced by another bent sequence of the same length. This replacing process is continued until the length of the all-one leading sequence is $2^2 = 4$. To finish the procedure, we replace the leading sequence $(1, 1, 1, 1)$ with $(1, -1, 1, -1)$. The last replacement makes the entire sequence balanced. By induction on $s = 2, 3, 4, \dots$, it can be proved that the nonlinearity of the function obtained is at least

$$2^{2^m-1} - \frac{1}{2}(2^{2^{m-1}} + 2^{2^{m-2}} + \dots + 2^{2^2} + 2 \cdot 2^2).$$

The modifying procedure for the case of $n = 2^s(2t + 1)$, $s \geq 1$ and $t \geq 1$, is the same as that for the case of $n = 2^m$, $m \geq 2$, except for the last replacement. In this case, the replacing process is continued until the length of the all-one leading sequence is 2^{2t+1} . The last leading sequence is replaced by $\ell_0^* = (e_{2^t}, e_{2^t+1}, \dots, e_{2^{2t+1}-1})$, the second half of the bent sequence $(e_0, e_1, \dots, e_{2^{2t+1}-1})$, where each e_i is a row of H_{t+1} . Again by induction on $s = 1, 2, 3, \dots$, it can be proved that the nonlinearity of the resulting function is at least

$$2^{2^s(2t+1)-1} - \frac{1}{2}(2^{2^{s-1}(2t+1)} + 2^{2^{s-2}(2t+1)} + \dots + 2^{2(2t+1)} + 2^{2t+1} + 2^{t+1}).$$

We have completed the proof for the following

Table 1: Nonlinearities of Balanced Functions on V_{2k}

Vector Space	V_4	V_6	V_8	V_{10}	V_{12}	V_{14}
Upper Bound (\leq)	4	26	118	494	2014	8126
By Modification (\geq)	4	26	116	492	2010	8120
By Concatenation (\geq)	4	24	112	480	1984	8064

Theorem 1 For any even number $n \geq 4$, there exists a balanced function f^* on V_n whose nonlinearity is

$$N_{f^*} \geq \begin{cases} 2^{2^m-1} - \frac{1}{2}(2^{2^m-1} + 2^{2^m-2} + \dots + 2^{2^2} + 2 \cdot 2^2), & n = 2^m, \\ 2^{2^s(2t+1)-1} - \frac{1}{2}(2^{2^{s-1}(2t+1)} + 2^{2^{s-2}(2t+1)} + \dots + 2^{2(2t+1)} + 2^{2t+1} + 2^{t+1}), & n = 2^s(2t+1). \end{cases}$$

Let $\zeta = (\zeta_0, \zeta_1, \dots, \zeta_{2^k-1})$ be a sequence of length 2^{2k} obtained by modifying a bent sequence. Permuting and changing signs discussed in Section 3.3 can also be applied to ζ . In this way we obtain in total $2^{2^k} \cdot 2^k!$ different balanced functions, all of which have the same nonlinearity. Even more functions can be obtained by observing the fact that the leading sequence ζ_0 has exactly the same structure as the large sequence ζ , and hence permuting and changing signs can also be applied to ζ_0 .

The nonlinearities of balanced functions on $V_4, V_6, V_8, V_{10}, V_{12}$ and V_{14} constructed by the method shown in the proof of Theorem 1 are calculated in Table 1. For comparison, the nonlinearities of balanced functions constructed by concatenating four bent sequences (see Lemma 8) as well as the upper bounds for the nonlinearities of balanced functions (see Lemma 5) are also presented.

4.2 On V_{2k+1}

The following lemma can be easily confirmed and very useful in obtaining balanced functions.

Lemma 12 Let ξ_1 be the sequence of f_1 on V_s and ξ_2 be the sequence of f_2 on V_t . Then

1. $f_1(x_1, \dots, x_s) \oplus f_2(y_1, \dots, y_t)$ is a balanced function on V_{s+t} if f_1 or f_2 is balanced.
2. The Kronecker product $\xi_1 \otimes \xi_2$ is the sequence of $f_1(x_1, \dots, x_s) \oplus f_2(y_1, \dots, y_t)$.

Lemma 13 Let ξ_1 be the sequence of f_1 on V_s and ξ_2 be the sequence of f_2 on V_t . Also let

$$g(x_1, \dots, x_s, y_1, \dots, y_t) = f_1(x_1, \dots, x_s) \oplus f_2(y_1, \dots, y_t).$$

Assume that $\langle \xi_1, \ell_1 \rangle \leq P_1$ and $\langle \xi_2, \ell_2 \rangle \leq P_2$, where ℓ_1 is an arbitrary affine sequence of length 2^s , ℓ_2 is an arbitrary affine sequence of length 2^t , P_1 and P_2 are positive integers. Then the nonlinearity of g satisfies $N_g \geq 2^{s+t-1} - \frac{1}{2}P_1 \cdot P_2$.

Proof. Note that $\xi = \xi_1 \otimes \xi_2$ is the sequence of g . Let φ be an arbitrary affine function on V_{s+t} and let ℓ be the sequence of φ . Then ℓ can be expressed as $\ell = \pm \ell_1 \otimes \ell_2$ where ℓ_1 is a row of H_s and ℓ_2 is a row of H_t . Since

$$\langle \xi, \ell \rangle = \langle \xi_1 \otimes \xi_2, \pm \ell_1 \otimes \ell_2 \rangle = \pm \langle \xi_1, \ell_1 \rangle \langle \xi_2, \ell_2 \rangle$$

we have

$$|\langle \xi, \ell \rangle| = |\langle \xi_1, \ell_1 \rangle| \cdot |\langle \xi_2, \ell_2 \rangle| \leq P_1 \cdot P_2$$

and by Lemma 6

$$d(g, \varphi) \geq 2^{s+t-1} - \frac{1}{2}P_1 \cdot P_2.$$

Due to the arbitrariness of φ , we have $N_g \geq 2^{s+t-1} - \frac{1}{2}P_1 \cdot P_2$. \square

Let ξ_1 be a balanced sequence of length 2^{2k} that is constructed using the method in the proof of Theorem 1, where $k \geq 2$, Let ξ_2 be a sequence of length 2^{15} obtained by the method of (Patterson & Wiedemann, 1983). Note that the nonlinearity of ξ_2 is 16276, and there are 13021 such sequences. Denote by f_1 the function corresponding to ξ_1 and by f_2 the function corresponding to ξ_2 . Let

$$f(x_1, \dots, x_{2k}, x_{2k+1}, \dots, x_{2k+15}) = f_1(x_1, \dots, x_{2k}) \oplus f_2(x_{2k+1}, \dots, x_{2k+15}). \quad (3)$$

Note that

$$\langle \xi_1, \ell_1 \rangle \leq \begin{cases} 2^{2^{m-1}} + 2^{2^{m-2}} + \dots + 2^{2^2} + 2 \cdot 2^2, & 2k = 2^m, \\ 2^{2^{s-1}(2t+1)} + 2^{2^{s-2}(2t+1)} + \dots + 2^{2^{2t+1}} + 2^{2t+1} + 2^{t+1}, & 2k = 2^s(2t+1). \end{cases}$$

and

$$\langle \xi_2, \ell_2 \rangle \leq 2 \cdot (2^{14} - 16276) = 216$$

where ℓ_1 is a linear sequence of length 2^{2k} and ℓ_2 is a linear sequence of length 2^{15} . Then by Lemma 13, we have:

Theorem 2 *The function f defined by (3) is a balanced function on V_{2k+15} , $k \geq 2$, whose nonlinearity is at least*

$$N_f \geq \begin{cases} 2^{2^m+14} - 108(2^{2^{m-1}} + 2^{2^{m-2}} + \dots + 2^{2^2} + 2 \cdot 2^2), & 2k = 2^m, \\ 2^{2^s(2t+1)+14} - 108(2^{2^{s-1}(2t+1)} + 2^{2^{s-2}(2t+1)} + \dots + 2^{2^{2t+1}} + 2^{2t+1} + 2^{t+1}), & 2k = 2^s(2t+1). \end{cases}$$

The nonlinearity of a function on V_{2k+15} constructed in this section is larger than that obtained by concatenating or splitting bent sequences for all $k \geq 7$.

5 Constructing Highly Nonlinear balanced Functions Satisfying SAC

This section presents methods for constructing balanced functions with a high nonlinearity and satisfying the SAC. The algebraic degrees of the functions are discussed.

5.1 On V_{2k+1}

Let $k \geq 1$, f a bent function and h a non-constant affine function, both on V_{2k} . Note that $f(x) \oplus h(x)$ is also bent. Without loss of generality we suppose that the number of times that $f(x)$ assumes the value zero differs from that of $f(x) \oplus h(x)$. Let g be a function on V_{2k+1} defined by

$$\begin{aligned} g(u, x_1, \dots, x_{2k}) &= (1 \oplus u)f(x_1, \dots, x_{2k}) \oplus u(f(x_1, \dots, x_{2k}) \oplus h(x_1, \dots, x_{2k})) \\ &= f(x_1, \dots, x_{2k}) \oplus uh(x_1, \dots, x_{2k}). \end{aligned} \quad (4)$$

By Lemma 2 the sequence of g is the concatenation of the sequences of $f(x)$ and $f(x) \oplus h(x)$. Thus we have

Lemma 14 *The function g defined by (4) is a balanced function on V_{2k+1} .*

The following lemma is a direct consequence of Corollary 1.

Lemma 15 *$N_g \geq 2^{2k} - 2^k$ where g is defined by (4).*

Lemma 16 *The function g defined by (4) satisfies the SAC.*

Proof. Let $\gamma = (b, a_1, \dots, a_{2k})$ be an arbitrary vector in V_{2k+1} with $W(\gamma) = 1$. Also let $\alpha = (a_1, \dots, a_{2k})$, $z = (u, x_1, \dots, x_{2k})$ and $x = (x_1, \dots, x_{2k})$. We show that $g(z) \oplus g(z \oplus \gamma) = f(x) \oplus f(x \oplus \alpha) \oplus u(h(x) \oplus h(x \oplus \alpha)) \oplus bh(x \oplus \alpha)$ is balanced by considering the following two cases.

Case 1: $b = 0$ and hence $W(\alpha) = 1$. Then $g(z) \oplus g(z \oplus \gamma) = f(x) \oplus f(x \oplus \alpha) \oplus u(h(x) \oplus h(x \oplus \alpha))$. Since h is an affine function, $h(x) \oplus h(x \oplus \alpha) = c$ where c is a constant from $GF(2)$. Thus $g(z) \oplus g(z \oplus \gamma) = f(x) \oplus f(x \oplus \alpha) \oplus cu$. By (iii) of Lemma 3, $f(x) \oplus f(x \oplus \alpha)$ is a balanced function on V_{2k} and hence by Lemma 12, $g(z) \oplus g(z \oplus \gamma)$ is a balanced function on V_{2k+1} .

Case 2: $b = 1$ and hence $W(\alpha) = 0$, i.e. $\alpha = (0, 0, \dots, 0)$. Then $g(z) \oplus g(z \oplus \gamma) = h(x)$. Since $h(x)$ is a non-constant affine function on V_{2k} , $h(x)$ and hence $g(z) \oplus g(z \oplus \gamma)$ are balanced. \square

Summarizing Lemmas 14, 15 and 16 we have

Theorem 3 *For $k \geq 1$, g defined by (4) is a balanced function on V_{2k+1} having $N_g \geq 2^{2k} - 2^k$ and satisfying the SAC.*

Recently Zheng, Pieprzyk and Seberry (1993) constructed a very efficient one way hashing algorithm using boolean functions constructed by the method given in Theorem 3. These functions have further cryptographically useful properties.

5.2 On V_{2k}

Let $k \geq 2$ and f a bent function on V_{2k-2} . And let h_1, h_2 and h_3 be non-constant affine functions on V_{2k-2} such that $h_i(x) \oplus h_j(x)$ is non-constant for any $i \neq j$. Such affine functions exist for all $k \geq 2$. Let $x = (x_1, \dots, x_{2k-2})$. Note that each $f(x) \oplus h_j(x)$ is also bent.

Without loss of generality we suppose both $f(x)$ and $f(x) \oplus h_1(x)$ assume the value one $2^{2k-3} + 2^{k-2}$ times while both $f(x) \oplus h_2(x)$ and $f(x) \oplus h_3(x)$ assume the value one $2^{2k-3} - 2^{k-2}$ times. Let g be a function on V_{2k} defined by

$$\begin{aligned} g(u, v, x_1, \dots, x_{2k-2}) &= (1 \oplus u)(1 \oplus v)f(x) \oplus (1 \oplus u)v(f(x) \oplus h_1(x)) \oplus \\ &\quad u(1 \oplus v)(f(x) \oplus h_2(x)) \oplus uv(f(x) \oplus h_3(x)) \\ &= f(x) \oplus vh_1(x) \oplus uh_2(x) \oplus uv(h_1(x) \oplus h_2(x) \oplus h_3(x)). \end{aligned} \quad (5)$$

Lemma 17 *g defined by (5) is a balanced function on V_{2k} .*

Proof. Note that the sequence of g is the concatenation of the sequences of $f(x)$, $f(x) \oplus h_1(x)$, $f(x) \oplus h_2(x)$ and $f(x) \oplus h_3(x)$, and that $f(x)$ and $f(x) \oplus h_1(x)$ assume the value one $2^{2k-3} + 2^{k-2}$ times while $f(x) \oplus h_2(x)$ and $f(x) \oplus h_3(x)$ assume the value one $2^{2k-3} - 2^{k-2}$ times. Thus g assumes the value one 2^{2k-1} times and hence is a balanced function on V_{2k} . \square

Lemma 18 $N_g \geq 2^{2k-1} - 2^k$ where g is defined by (5).

Proof. It follows from Corollary 1. □

Lemma 19 The function g defined by (5) satisfies the SAC.

Proof. Let $\gamma = (b, c, a_1, \dots, a_{2k-2})$ be any vector in V_{2k} with $W(\gamma) = 1$. Write $\alpha = (a_1, \dots, a_{2k-2})$, $z = (u, v, x_1, \dots, x_{2k-2})$ and $x = (x_1, \dots, x_{2k-2})$. Note that $g(z \oplus \gamma) = f(x \oplus \alpha) \oplus (v \oplus c)h_1(x \oplus \alpha) \oplus (u \oplus b)h_2(x \oplus \alpha) \oplus (u \oplus b)(v \oplus c)(h_1(x \oplus \alpha) \oplus h_2(x \oplus \alpha) \oplus h_3(x \oplus \alpha))$. Consider the balance of $g(z) \oplus g(z \oplus \gamma)$ in the following three cases.

Case 1: $b = 1, c = 0$ and hence $W(\alpha) = 0$, i.e. $\alpha = (0, 0, \dots, 0)$. In this case, $g(z) \oplus g(z \oplus \gamma) = h_2(x) \oplus v(h_1(x) \oplus h_2(x) \oplus h_3(x))$ will be $h_2(x)$ when $v = 0$ and $h_1(x) \oplus h_3(x)$ when $v = 1$. Both $h_2(x)$ and $h_1(x) \oplus h_3(x)$ are non-constant affine functions on V_{2k-2} and hence $g(z) \oplus g(z \oplus \gamma)$ is a balanced function on V_{2k} .

Case 2: $b = 0, c = 1$ and hence $W(\alpha) = 0$, i.e. $\alpha = (0, 0, \dots, 0)$. The proof of the balance of $g(z) \oplus g(z \oplus \gamma)$ is similar to Case 1.

Case 3: $b = 0, c = 0$ and hence $W(\alpha) = 1$. Since h_j is an affine function, $h_j(x) \oplus h_j(x \oplus \alpha) = a_j$ where a_j is a constant from $GF(2)$. Hence $g(z) \oplus g(z \oplus \gamma) = f(x) \oplus f(x \oplus \alpha) \oplus va_1 \oplus ua_2 \oplus uv(a_1 \oplus a_2 \oplus a_3)$. By (iii) of Lemma 3, $f(x) \oplus f(x \oplus \alpha)$ is a balanced function on V_{2k-2} and hence by Lemma 12, $g(z) \oplus g(z \oplus \gamma)$ is a balanced function on V_{2k} . This proves that g satisfies the SAC. □

Summarizing Lemmas 17, 18 and 19 we have

Theorem 4 For $k \geq 2$, g defined by (5) is a balanced function on V_{2k} having $N_g \geq 2^{2k-1} - 2^k$ and satisfying the SAC.

5.3 Remarks

We have shown that a function on V_n constructed according to (4) and (5) satisfy the propagation criterion with respect to all the n vectors whose Hamming weight is 1. In fact there are many more vectors where the propagation criterion is satisfied, and it is not hard to show that the total number of vectors in V_{2k+1} such that a function constructed by (4) satisfies the propagation criterion is $2^{2k} + 2^{2k-1}$, while the total number of vectors in V_{2k} such that a function constructed by (5), satisfies the propagation criterion is at least $2^{2k-2} + 1$.

The algebraic degree is also a nonlinearity criterion and it becomes important in certain practical applications where linear approximation of a nonlinear function needs to be avoided. In our constructions (4) and (5), the algebraic degree of a resulting function g is the same as that of the starting bent function f .

The simplest bent function on V_{2k} is the following quadratic function:

$$f(x_1, x_2, \dots, x_{2k}) = x_1x_{k+1} \oplus x_2x_{k+2} \oplus \dots \oplus x_kx_{2k}.$$

Bent functions with higher algebraic degrees exist and there are many methods for constructing such functions (Dillon, 1972). The following is a method discovered by Dillon and Maiorana (1972; 1985) for constructing a bent function f on V_{2k} :

$$f(x) = \langle x', \pi(x'') \rangle \oplus r(x'')$$

where $x = (x', x'')$, $x' = (x_1, \dots, x_k)$, $x'' = (x_{k+1}, \dots, x_{2k})$, r is an arbitrary function on V_k and $\pi = (\pi_1(x''), \pi_2(x''), \dots, \pi_k(x''))$ is a permutation on the vector space V_k . Due to the arbitrariness of r , the algebraic degree of f can be any integer between 2 and k . From these discussions it becomes clear that functions obtained by (4) and (5) can achieve a wide range of algebraic degrees, namely $2, \dots, k$ and $2, \dots, k - 1$ respectively.

6 Constructing Highly Nonlinear balanced Functions Satisfying High Degree Propagation Criterion

Another interesting topic is to study methods for constructing functions that are balanced and possess good propagation characteristics. In (Preneel *et al.*, 1991a), it was suggested that a function f on V_n which has a zero point in its Walsh spectrum be modified into a balanced function by adding a suitable linear function h on V_n . As h has to be found by exhaustive search over all the linear functions on V_n , the method is infeasible when n is large. In addition, the method is not applicable to the functions which do not have zero points in their Walsh spectra. Two types of such functions are (1) bent functions, and (2) highly nonlinear functions obtained by complementing a single position in a bent sequence. In the following we describe two methods for systematically constructing highly nonlinear balanced functions satisfying high degree propagation criterion.

6.1 Basic Construction

6.1.1 On V_{2k+1}

Let f be a bent function on V_{2k} , and let g be a function on V_{2k+1} defined by

$$\begin{aligned} g(x_1, x_2, \dots, x_{2k+1}) &= (1 \oplus x_1)f(x_2, \dots, x_{2k+1}) \oplus x_1(1 \oplus f(x_2, \dots, x_{2k+1})) \\ &= x_1 \oplus f(x_2, \dots, x_{2k+1}). \end{aligned} \quad (6)$$

Lemma 20 *The function g defined in (6) satisfies the propagation criterion with respect to all non-zero vectors $\gamma \in V_{2k+1}$ with $\gamma \neq (1, 0, \dots, 0)$.*

Proof. Let $\gamma = (a_1, a_2, \dots, a_{2k+1}) \neq (1, 0, \dots, 0)$ and let $x = (x_1, x_2, \dots, x_{2k+1})$. Then $g(x) \oplus g(x \oplus \gamma) = a_1 \oplus f(x_2, \dots, x_{2k+1}) \oplus f(x_2 \oplus a_2, \dots, x_{2k+1} \oplus a_{2k+1})$. Since f is a bent function, $f(x_2, \dots, x_{2k+1}) \oplus f(x_2 \oplus a_2, \dots, x_{2k+1} \oplus a_{2k+1})$ is balanced for all $(a_2, \dots, a_{2k+1}) \neq (0, \dots, 0)$ (see (iii) of Lemma 3). Thus $g(x) \oplus g(x \oplus \gamma)$ is balanced for all $\gamma = (a_1, a_2, \dots, a_{2k+1}) \neq (1, 0, \dots, 0)$. \square

From Corollary 1, the nonlinearity of the function g defined by (6) satisfies $N_g \geq 2^{2k} - 2^k$. Furthermore, by Lemma 12, g is balanced. Thus we have

Corollary 2 *The function g defined by (6) is balanced and satisfies the propagation criterion with respect to all non-zero vectors $\gamma \in V_{2k+1}$ with $\gamma \neq (1, 0, \dots, 0)$. The nonlinearity of g satisfies $N_g \geq 2^{2k} - 2^k$.*

6.1.2 On V_{2k}

Let f be a bent function on V_{2k-2} and let g be a function on V_{2k} obtained from f in the following way:

$$g(x_1, x_2, x_3, \dots, x_{2k})$$

$$\begin{aligned}
&= (1 \oplus x_1)(1 \oplus x_2)f(x_3, \dots, x_{2k}) \oplus (1 \oplus x_1)x_2(1 \oplus f(x_3, \dots, x_{2k})) \\
&\quad x_1(1 \oplus x_2)(1 \oplus f(x_3, \dots, x_{2k})) \oplus x_1x_2f(x_3, \dots, x_{2k}) \\
&= x_1 \oplus x_2 \oplus f(x_3, \dots, x_{2k}). \tag{7}
\end{aligned}$$

Lemma 21 *The function g defined in (7) satisfies the propagation criterion with respect to all but three non-zero vectors in V_{2k} . The three vectors where the propagation criterion is not satisfied are $\gamma_1 = (1, 0, 0, \dots, 0)$, $\gamma_2 = (0, 1, 0, \dots, 0)$, and $\gamma_3 = \gamma_1 \oplus \gamma_2 = (1, 1, 0, \dots, 0)$.*

Proof. Let $\gamma = (a_1, a_2, \dots, a_{2k})$ be a non-zero vector in V_{2k} differing from γ_1, γ_2 and γ_3 . Also let $x = (x_1, \dots, x_{2k})$. Then we have $g(x) \oplus g(x \oplus \gamma) = a_1 \oplus a_2 \oplus f(x_3, \dots, x_{2k}) \oplus f(x_3 \oplus a_3, \dots, x_{2k} \oplus a_{2k})$. Since f is a bent function on V_{2k-2} and $(a_3, \dots, a_{2k}) \neq (0, \dots, 0)$, $f(x_3, \dots, x_{2k}) \oplus f(x_3 \oplus a_3, \dots, x_{2k} \oplus a_{2k})$ is balanced, from which it follows that $g(x) \oplus g(x \oplus \gamma)$ is balanced for any non-zero vector γ in V_{2k} differing from γ_1, γ_2 and γ_3 . This proves the lemma. \square

Since $x_1 \oplus x_2$ is balanced on V_2 , g is balanced on V_{2k} . On the other hand, by Lemma 7, we have $N_g \geq 2^{2k-1} - 2^k$. Thus we have the following result:

Corollary 3 *The function g defined by (7) is balanced and satisfies the propagation criterion with respect to all non-zero vectors $\gamma \in V_{2k}$ with $\gamma \neq (c_1, c_2, 0, \dots, 0)$, where $c_1, c_2 \in GF(2)$. The nonlinearity of g satisfies $N_g \geq 2^{2k-1} - 2^k$.*

6.2 Moving Vectors Around

Though functions constructed according to (6) or (7) satisfy the propagation criterion with respect to all but one or three non-zero vectors, they only fulfill the propagation criterion of degree zero. Therefore these functions are not interesting in practical applications. Recall that the balance, the nonlinearity and the number of vectors where the propagation criterion is satisfied are all invariant under an affine transformation of coordinates. This indicates that the degree for the propagation criterion might be improved through a suitable affine transformation of coordinates. Identifying such an affine transformation, however, is not an easy exercise, especially when the dimension of the underlying vector space is large and the number of vectors where the propagation criterion is satisfied is small.

In this section, we show that for functions constructed according to (6) or (7), the vectors where the propagation criterion is not satisfied can be transformed into vectors having a high Hamming weight. In this way we obtain highly nonlinear balanced functions satisfying high degree propagation criterion.

6.2.1 On V_{2k+1}

Theorem 5 *For any non-zero vector $\gamma^* \in V_{2k+1}$ ($k \geq 1$), there exist balanced functions on V_{2k+1} satisfying the propagation criterion with respect to all non-zero vectors $\gamma \in V_{2k+1}$ with $\gamma \neq \gamma^*$. The nonlinearities of the functions are at least $2^{2k} - 2^k$.*

Proof. Let f be a bent function and let g be the function constructed by (6). From linear algebra we know that for any bases B_1 and B_2 of the vector space V_{2k+1} , where $B_1 = \{\alpha_j | j = 1, \dots, 2k+1\}$ and $B_2 = \{\beta_j | j = 1, \dots, 2k+1\}$, there exists a unique nondegenerate matrix A of order $2k+1$ with entries from $GF(2)$ such that $\alpha_j A = \beta_j$, $j = 1, \dots, 2k+1$. In particular, this is true when

$\alpha_1 = \gamma^*$ and $\beta_1 = (1, 0, \dots, 0)$. Let $x = (x_1, x_2, \dots, x_n)$ and let g^* be the function obtained from g by employing linear transformation on the input coordinates of g :

$$g^*(x) = g(xA).$$

Since A is nondegenerate, by Lemma 10, g^* is balanced and has the same nonlinearity as that of g . Now we show that g^* satisfies the propagation criterion with respect to all non-zero vectors except γ^* .

Let γ be a non-zero vector in V_{2k+1} with $\gamma \neq \gamma^*$. Consider the following function $g^*(x) \oplus g^*(x \oplus \gamma) = g(xA) \oplus g(xA \oplus \gamma A) = g(y) \oplus g(y \oplus \gamma A)$ where $y = xA$. Note that A is nondegenerate and thus y runs through V_{2k+1} while x runs through V_{2k+1} . Since $\gamma \neq \gamma^*$ we have $\gamma A \neq (1, 0, \dots, 0)$. By Lemma 20, $g(y) \oplus g(y \oplus \gamma A)$ runs through the values zero and one an equal number of times. Hence $g^*(x) \oplus g^*(x \oplus \gamma)$ is balanced. Consequently, g^* satisfies the propagation criterion with respect to all non-zero vectors in V_{2k+1} but γ^* . This completes the proof. \square

As a consequence of Theorem 5, we obtain, by letting $\gamma^* = (1, 1, \dots, 1)$, highly nonlinear balanced functions on V_{2k+1} satisfying the propagation criterion of degree $2k$. This is described in the following:

Corollary 4 *Let f be a bent function on V_{2k} and let $g^*(x_1, \dots, x_{2k+1}) = x_1 \oplus f(x_1 \oplus x_2, x_1 \oplus x_3, \dots, x_1 \oplus x_{2k+1})$. Then g^* is a balanced function on V_{2k+1} and satisfies the propagation criterion of degree $2k$. The nonlinearity of g^* satisfies $N_{g^*} \geq 2^{2k} - 2^k$.*

Proof. Let e_j , $j = 1, 2, \dots, 2k+1$, be a vector in V_{2k+1} whose j th coordinate is 1 and all other coordinates are 0. In the proof of Theorem 5, we let $\alpha_1 = \gamma_0 = (1, \dots, 1)$, $\alpha_j = e_j$, $j = 2, \dots, 2k+1$ and $\beta_j = e_j$, $j = 1, \dots, 2k+1$. Then there is a unique nondegenerate matrix A of order $2k+1$ such that $\alpha_j A = \beta_j$, $j = 1, \dots, 2k+1$. It is easy to verify that A has the following form:

$$A = \begin{bmatrix} \gamma_0 \\ e_2 \\ \vdots \\ e_{2k+1} \end{bmatrix}.$$

Thus we have $g^*(x) = g(xA) = g(x_1, x_1 \oplus x_2, \dots, x_1 \oplus x_{2k+1}) = x_1 \oplus f(x_1 \oplus x_2, x_1 \oplus x_3, \dots, x_1 \oplus x_{2k+1})$, where $g(x) = x_1 \oplus f(x_2, \dots, x_{2k+1})$, and $x = (x_1, x_2, \dots, x_{2k+1})$. By Theorem 5 g^* satisfies the propagation criterion with respect to all non-zero vectors in V_{2k+1} except the all-one vector $\gamma^* = (1, 1, \dots, 1)$. Consequently g^* satisfies the propagation criterion of degree $2k$. \square

6.2.2 On V_{2k}

Theorem 6 *For any non-zero vectors $\gamma_1^*, \gamma_2^* \in V_{2k}$ ($k \geq 2$) with $\gamma_1^* \neq \gamma_2^*$, there exist balanced functions on V_{2k} satisfying the propagation criterion with respect to all but three non-zero vectors in V_{2k} . The three vectors where the propagation criterion is not satisfied are γ_1^* , γ_2^* and $\gamma_1^* \oplus \gamma_2^*$. The nonlinearities of the functions are at least $2^{2k-1} - 2^k$.*

Proof. The proof is essentially the same as that for Theorem 5. The major difference lies in the selection of bases $B_1 = \{\alpha_j | j = 1, \dots, 2k\}$ and $B_2 = \{\beta_j | j = 1, \dots, 2k\}$. By linear algebra, we can let $\alpha_1 = \gamma_1^*$, $\alpha_2 = \gamma_2^*$, $\beta_1 = (1, 0, 0, \dots, 0)$, and $\beta_2 = (0, 1, 0, \dots, 0)$. By the same reasoning as

in the proof of Theorem 5, we can see that g^* defined by $g^*(x) = g(xA)$ satisfies the propagation criterion with respect to all but the following three non-zero vectors in V_{2k} : γ_1^* , γ_2^* and $\gamma_1^* \oplus \gamma_2^*$. Here $x = (x_1, x_2, \dots, x_{2k})$, $g(x) = x_1 \oplus x_2 \oplus f(x_3, \dots, x_{2k})$, and f , a bent function on V_{2k-2} , are all the same as in (7), and A is the unique nondegenerate matrix such that $\alpha_j A = \beta_j$, $j = 1, \dots, 2k$. \square

Similarly to the case on V_{2k+1} , we can obtain highly nonlinear balanced functions satisfying high degree propagation criterion, by properly selecting vectors γ_1^* and γ_2^* . Unlike the case on V_{2k+1} , however, the degree of propagation criterion the functions can achieve is $\frac{4}{3}k$, but not $2k - 1$. The construction method is described in the following corollary.

Corollary 5 *Suppose that $2k = 3t + c$ where $c = 0, 1$ or 2 . Then there exist balanced functions on V_{2k} that satisfy the propagation criterion of degree $2t - 1$ (when $c = 0$ or 1), or $2t$ (when $c = 2$). The nonlinearities of the functions are at least $2^{2k-1} - 2^k$.*

Proof. Set $c_1 = 0$, $c_2 = 1$ if $c = 1$ and set $c_1 = c_2 = \frac{1}{2}c$ otherwise. Let $\gamma_1^* = (a_1, \dots, a_{3t+c})$ and $\gamma_2^* = (b_1, \dots, b_{3t+c})$, where

$$a_j = \begin{cases} 1 & \text{for } j = 1, \dots, 2t + c_1, \\ 0 & \text{for } j = 2t + c_1 + 1, \dots, 3t + c. \end{cases}$$

$$b_j = \begin{cases} 0 & \text{for } j = 1, \dots, t + c_1, \\ 1 & \text{for } j = t + c_1 + 1, \dots, 3t + c. \end{cases}$$

By Theorem 6 there exists a balanced function g^* on V_{2k} satisfying the propagation criterion with respect to all but three non-zero vectors in V_{2k} . The three vectors are γ_1^* , γ_2^* and $\gamma_1^* \oplus \gamma_2^*$. The nonlinearity of g^* satisfies $N_{g^*} \geq 2^{2k-1} - 2^k$.

Note that $W(\gamma_1^*) = 2t + c_1$, $W(\gamma_2^*) = 2t + c_2$, and $W(\gamma_1^* \oplus \gamma_2^*) = 2t + 2c_1 = 2t + c$. The minimum among the three weights is $2t + c_1$. Therefore, for any nonzero vector $\gamma \in V_{2k}$ with $W(\gamma) \leq 2t + c_1 - 1$, we have $\gamma \neq \gamma_1^*, \gamma_2^*$ or $\gamma_1^* \oplus \gamma_2^*$. By Theorem 6, $g^*(x) \oplus g^*(x \oplus \gamma)$ is balanced. From this we conclude that g^* satisfies the propagation criterion of order $2t + c_1 - 1$. The proof is completed by noting that $c_1 = 0$ if $c = 0$ or 1 and $c_1 = 1$ if $c = 2$. \square

6.3 Discussions

Comparing (4) with (6), one can see that the difference between the two constructions lies in the selection of the affine functions. In (4) a *non-constant* affine function h is selected, while in (6) a constant 1 is employed. In a sense, the two constructions complement one another. A similar observation applies to the case of (5) and (7).

Functions obtained by (6) and (7) can achieve a wide range of algebraic degrees, namely $2, \dots, k$ and $2, \dots, k - 1$ respectively. (See also the discussions in Section 5.3.) Recently, Detombe and Tavares obtained, while studying the design of S-boxes, balanced *quadratic* functions on V_5 that satisfy the propagation criterion with respect to all but one vectors in V_5 . (They called these functions *near bent* functions.) They obtained the functions by the use of the *cubing* technique suggested by Pieprzyk (1991). Propagation characteristics of quadratic functions were also studied extensively in (Preneel *et al.*, 1991a). However, applicability of these quadratic functions in practice is limited by the following two facts:

1. Their algebraic degree is only 2.
2. They are all equivalent in structure in the sense that they can be transformed into one another by linear transformation of input coordinates.

7 Concluding Remarks

We have studied properties of balance and nonlinearity of Boolean functions including concatenating, splitting, modifying and multiplying sequences. A novel method has been presented to construct balanced functions whose nonlinearity is much higher than that attained by any previously known construction. In addition, systematic methods have been presented for constructing highly nonlinear balanced functions satisfying the SAC or high degree propagation criterion. A technique has been developed that allows us to transform vectors where the propagation criterion is not satisfied into other vectors, while preserving the nonlinearity and balance of the functions. This paper has also introduced a number of interesting problems which remain to be solved. We discuss one of them before closing the paper. For V_{2k+1} , functions constructed according to (6) are optimal in the sense that they fulfill the propagation criterion with respect to $2^{2k+1} - 2$ non-zero vectors, and after the affine transformation of coordinates, they satisfy the propagation criterion of degree $2k$. For V_{2k} , the number of non-zero vectors given by (7) is $2^{2k} - 4$ and the degree after the transformation is $\frac{4k}{3}$. It is left as future work to examine whether there are highly nonlinear balanced functions on V_{2k} satisfying the propagation criterion of degree $2k - 1$, and if there are, to find methods for constructing such functions.

Acknowledgments

This work was supported by Telecom Australia under the contract number 7027 and by the Australian Research Council under the reference numbers A48830241, A49130102, A9030136, A49131885 and A49232172. The authors would like to thank the anonymous referees for their helpful comments.

References

- ADAMS, C. M., & TAVARES, S. E. 1990a. Generating and Counting binary bent sequences. *IEEE Transactions on Information Theory*, **IT-36 No. 5**, 1170–1173.
- ADAMS, C. M., & TAVARES, S. E. 1990b. *The use of bent sequences to achieve higher-order strict avalanche criterion*. Technical Report, TR 90-013, Department of Electrical Engineering, Queen's University.
- COHEN, G. D., KARPOVSKY, M. G., H. F. MATTSON, JR., & SCHATZ, J. R. 1985. Covering Radius — Survey and Recent Results. *IEEE Transactions on Information Theory*, **IT-31(3)**, 328–343.
- DETOMBE, J., & TAVARES, S. 1993. Constructing Large Cryptographically Strong S-boxes. *Pages 165–181 of: Advances in Cryptology - AUSCRYPT'92*, vol. 718, Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, New York.
- DILLON, J. F. 1972. A Survey of Bent Functions. *The NSA Technical Journal*, 191–215. (unclassified).
- FORRÉ, R. 1989. The Strict Avalanche Criterion: Special Properties of Boolean Functions and Extended Definition. *Pages 450–468 of: Advances in Cryptology - CRYPTO'88*, vol. 403, Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, New York.

- KAM, J. B., & DAVIDA, G. I. 1979. Structured Design of Substitution-Permutation Encryption Networks. *IEEE Transactions on Computers*, **28**, 747–753.
- KUMAR, P. V., & SCHOLTZ, R. A. 1983. Bounds on the linear span of bent sequences. *IEEE Transactions on Information Theory*, **IT-29 No. 6**, 854–862.
- KUMAR, P. V., SCHOLTZ, R. A., & WELCH, L. R. 1985. Generalized Bent Functions and their properties. *Journal of Combinatorial Theory*, **Ser. A, 40**, 90–107.
- LEMPEL, A., & COHN, M. 1982. Maximal families of bent sequences. *IEEE Transactions on Information Theory*, **IT-28 No. 6**, 865–868.
- LOSEV, V. V. 1987. Decoding of sequences of bent functions by means of a fast Hadamard transform. *Radiotekhnika i elektronika*, **7**, 1479–1492.
- MACWILLIAMS, F. J., & SLOANE, N. J. A. 1978. *The Theory of Error-Correcting Codes*. Amsterdam, New York, Oxford: North-Holland.
- MATSUI, M. 1994. Linear Cryptanalysis Method for DES Cipher. *Pages 386–397 of: Advances in Cryptology - EUROCRYPT'93*, vol. 765, Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, New York.
- MEIER, W., & STAFFELBACH, O. 1990. Nonlinearity Criteria For Cryptographic Functions. *Pages 549–562 of: Advances in Cryptology - EUROCRYPT'89*, vol. 434, Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, New York.
- NYBERG, K. 1991. Perfect Nonlinear S-boxes. *Pages 378–386 of: Advances in Cryptology - EUROCRYPT'91*, vol. 547, Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, New York.
- OLSEN, J. D., SCHOLTZ, R. A., & WELCH, L. R. 1982. Bent-function sequences. *IEEE Transactions on Information Theory*, **IT-28 No. 6**, 858–864.
- PATTERSON, N. J., & WIEDEMANN, D. H. 1983. The Covering Radius of the $(2^{15}, 16)$ Reed-Muller Code Is at Least 16276. *IEEE Transactions on Information Theory*, **IT-29(3)**, 354–356.
- PIEPRZYK, J. 1991. Bent permutations. *In: Proceeding of the International Conference on Finite Fields, Coding Theory, and Advances in Communications and Computing*.
- PRENEEL, B., GOVAERTS, R., & VANDEWALLE, J. 1991a. Boolean Functions Satisfying Higher Order Propagation Criteria. *Pages 141–152 of: Advances in Cryptology - EUROCRYPT'91*, vol. 547, Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, New York.
- PRENEEL, B., LEEKWIJCK, W. V., LINDEN, L. V., GOVAERTS, R., & VANDEWALLE, J. 1991b. Propagation Characteristics of Boolean Functions. *Pages 155–165 of: Advances in Cryptology - EUROCRYPT'90*, vol. 437, Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, New York.
- ROTHAUS, O. S. 1976. On “Bent” Functions. *Journal of Combinatorial Theory*, **Ser. A, 20**, 300–305.
- SEBERRY, J., ZHANG, X. M., & ZHENG, Y. 1993. Systematic Generation of Cryptographically Robust S-boxes. *Pages 172 – 182 of: Proceedings of the first ACM Conference on Computer and Communications Security*. The Association for Computing Machinery, New York.

- WALLIS, W. D., STREET, A. PENFOLD, & WALLIS, J. SEBERRY. 1972. *Combinatorics: Room Squares, sum-free sets, Hadamard Matrices. Lecture Notes in Mathematics*, vol. 292. Springer-Verlag, Berlin, Heidelberg, New York.
- WEBSTER, A. F. 1985. *Plaintext/Ciphertext Bit Dependencies in Cryptographic System*. Master's Thesis, Department of Electrical Engineering, Queen's University, Ontario.
- WEBSTER, A. F., & TAVARES, S. E. 1986. On the Design of S-Boxes. *Pages 523–534 of: Advances in Cryptology - CRYPTO'85*, vol. 219, Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, New York.
- YARLAGADDA, R., & HERSHEY, J. E. 1989. Analysis and synthesis of bent sequences. *IEE Proceedings (Part E)*, **136**, 112–123.
- ZHENG, Y., PIEPRZYK, J., & SEBERRY, J. 1993. HAVAL - One-Way Hashing Algorithm with Variable Length of Output. *Pages 83–104 of: Advances in Cryptology - AUSCRYPT'92*, vol. 718, Lecture Notes in Computer Science. Springer-Verlag, Berlin, Heidelberg, New York.